

Northern Michigan University

NMU Commons

Journal Articles

FacWorks

2-2014

Midbrain Volume Predicts fMRI and ERP Measures of Reward Reactivity

Joshua M. Carlson
joshcarl@nmu.edu

Dan Foti

Eddie Harmon-Jones

Greg H. Proudfit

Follow this and additional works at: https://commons.nmu.edu/facwork_journalarticles

Recommended Citation

Carlson, J. M., Foti, D., Harmon-Jones, E., & Proudfit, G. H. (2014). Midbrain volume predicts fMRI and ERP measures of reward reactivity. *Brain Structure and Function*, 220(3), 1861-1866. <http://dx.doi.org/10.1007/s00429-014-0725-9>

This Journal Article is brought to you for free and open access by the FacWorks at NMU Commons. It has been accepted for inclusion in Journal Articles by an authorized administrator of NMU Commons. For more information, please contact kmcdonou@nmu.edu, bsarjean@nmu.edu.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/260253340>

Midbrain volume predicts fMRI and ERP measures of reward reactivity

Article in *Brain Structure and Function* · February 2014

DOI: 10.1007/s00429-014-0725-9 · Source: PubMed

CITATIONS

13

READS

90

4 authors, including:



Joshua M Carlson

Northern Michigan University

49 PUBLICATIONS 706 CITATIONS

SEE PROFILE



Dan Foti

Purdue University

55 PUBLICATIONS 1,701 CITATIONS

SEE PROFILE



Eddie Harmon-Jones

UNSW Australia

211 PUBLICATIONS 10,692 CITATIONS

SEE PROFILE

Midbrain volume predicts fMRI and ERP measures of reward reactivity

Joshua M. Carlson · Dan Foti · Eddie Harmon-Jones · Greg H. Proudfit

Received: 24 October 2013 / Accepted: 29 January 2014 / Published online: 19 February 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract Ventral striatal activation measured with functional magnetic resonance imaging (fMRI) and feedback negativity amplitude measured with event-related potentials (ERPs) are each enhanced during reward processing. Recent research has found that these two neural measures of reward processing are also related to one another, such that increases in ventral striatal activity are accompanied by increases in the amplitude of the feedback negativity. Although there is a long history of research implicating the midbrain dopamine system in reward processing, there has been little research into the possibility that structural variability in the midbrain may be linked to functional variability in reward reactivity. Here, we used structural MRI to measure midbrain volumes in addition to fMRI and ERP measures of functional neural reactivity to rewards in a simple gambling task. The results suggest that as midbrain volumes increase, fMRI reward reactivity in the ventral striatum and medial prefrontal cortex also increases. A similar relationship exists between midbrain structure and the amplitude of the feedback negativity; further, this

relationship is mediated specifically by activity in the ventral striatum. These data demonstrate convergence between neuroanatomical, hemodynamic, and electrophysiological measures. Thus, structural variability in the midbrain relates to variability in fMRI and ERP measures of functional reward reactivity, which may play a critical role in reward-related psychopathologies and the treatment of these disorders.

Keywords Reward · Striatum · Volume · Morphology · Feedback negativity · VBM

Introduction

The ability to crave, seek, detect, obtain, and enjoy resources or actions that benefit an organism's survival is an essential aspect of motivated drive states. A long history of neuroscience research in animals has implicated the dopaminergic projections from the midbrain to the ventral striatum (VS) in reward processing (Olds and Milner 1954; Wise 1996). In human functional magnetic resonance imaging (fMRI) studies the VS activates in response to a number rewarding stimuli such as drugs (Breiter et al. 1997; Drevets et al. 2001), erotic images (Walter et al. 2008; Sabatinelli et al. 2007), pleasant tastes (O'Doherty et al. 2002), attractive faces (Senior 2003), monetary rewards (Knutson and Bossaerts 2007), and favorable social interactions (Zink et al. 2008). Additional evidence using event-related potentials (ERPs) has identified the "feedback negativity" (FN; 300 ms) as an evoked potential that is sensitive to rewarding versus non-rewarding stimuli (Gehring and Willoughby 2002; Miltner et al. 1997; Holroyd et al. 2008; Foti et al. 2011). The amplitude of the FN is thought to track the relative valence of outcomes

J. M. Carlson (✉)
Department of Psychology, Northern Michigan University,
Marquette, MI 49855, USA
e-mail: joshcarl@nmu.edu

D. Foti
Department of Psychology, Purdue University, West Lafayette,
IN, USA

E. Harmon-Jones
School of Psychology, The University of New South Wales,
Sydney, Australia

G. H. Proudfit
Department of Psychology, Stony Brook University,
Stony Brook, NY, USA

(Holroyd et al. 2006, 2004). Recent research has explored the relationship between fMRI and EEG measures of reward anticipation (Plichta et al. 2013) and reward feedback (Carlson et al. 2011). In particular, research by our group demonstrates that increased VS activity is linked to increased FN amplitude (Carlson et al. 2011).

Ventral Striatal activity (Wise 1996) and FN amplitude (Holroyd and Coles 2002) are both thought to be influenced by the dopaminergic midbrain. Yet, the extent to which structural variability in midbrain morphology relates to variability in these measures of neural reactivity remains unknown. The importance of establishing the relationship between brain structure and reward reactivity is highlighted by research linking blunted reward processing to depressive symptoms. Indeed, both blunted VS activity (Epstein et al. 2006; Fitzgerald et al. 2008; Pizzagalli et al. 2009) and FN amplitude (Foti and Hajcak 2009) to rewards have been linked to heightened levels of depression. In addition, research suggests that a blunted FN amplitude is associated with depression as early as late childhood (Bress et al. 2012, 2013b) and a blunted FN prospectively predicts the onset of a major depressive episode (Bress et al. 2013a). The symptom of anhedonia in particular has been linked to blunted VS reward reactivity (Keedwell et al. 2005; Wacker et al. 2009) and deep brain stimulation of the VS attenuates the symptoms of anhedonia in depressed individuals (Schlaepfer et al. 2008). It is possible that these individual differences in reward reactivity may be driven in part by structural differences in midbrain morphology. Thus, given the importance of reward processing in normative adaptive behavior and depressive behavior, it is essential to understand the structure–function relationship of reward processing in the brain.

The focus of the current study was to link functional measures of reward processing as measured by fMRI and ERP (i.e., VS activation and FN amplitude, respectively) to underlying variability in neural structure within the dopaminergic midbrain as measured by structural MRI. Participants completed a gambling task, which contained equal numbers of trials containing monetary wins and losses in both fMRI and ERP environments (Foti and Hajcak 2009; Hajcak et al. 2006). Given the longstanding association between midbrain-regulated dopaminergic neurotransmission and reward processing, we predicted that the magnitude of reward-related reactivity in the VS and FN across individuals would positively correlate with variability in midbrain volume.

Methods

Participants

The same sample of 45 (Male = 27) consenting adults between the ages of 19 and 25 ($M = 21.11$, $SD = 1.27$)

that participated in our previously published study (Carlson et al. 2011) assessing correlations between fMRI and ERP measures was also used for the current manuscript. Participants were screened for metal and were monetarily compensated for their time. The Institutional Review Board of Stony Brook University approved this study.

Gambling task

The details of the experimental task as well as the fMRI and EEG preprocessing and analysis procedures have previously been described in detail (Carlson et al. 2011). Functional MRI and EEG data were collected in separate counterbalanced sessions, while participants performed a simple gambling task. Briefly, trials began with a white fixation cue presented in the center of a black screen. After which, a screen displayed two doors side-by-side and participants were instructed to choose a door. Behind one of the doors there was a monetary prize (+\$0.50), while behind the other door there was a loss (−\$0.25). After another brief fixation cue, a feedback screen was displayed where a green ‘↑’ indicated a win, and a red ‘↓’ indicated a loss. The task consisted of 60 trials with 30 predetermined wins and losses presented in a pseudorandom order.

EEG and fMRI measures

Win > Loss VS activity and FN amplitude data were used in a previous report (Carlson et al. 2011). Here, we utilized the same extracted BOLD data from the left and right VS as well as the medial prefrontal cortex (mPFC; left hemisphere maxima extending bilaterally). In addition, *Win > Loss* ERP activity corresponding to the FN was extracted using temporal-spatial principal components analysis (FN_{PCA}). The details regarding these measures can be obtained in our earlier report (Carlson et al. 2011). As previously reported, the FN_{PCA} correlated with the VS (left: $r = 0.28$, $p < 0.05$ and right: $r = 0.52$, $p < 0.001$) and mPFC ($r = 0.26$, $p < 0.05$) as well as other reward-related regions (Carlson et al. 2011).

Structural image acquisition and analysis

A 3 Tesla Siemens Trio whole body scanner was used to acquire T₁ images with the following parameters: TR = 1,900 ms, TE = 2.53, flip angle = 9°, FOV = 176 × 250 × 250 mm, matrix = 176 × 256 × 256, and voxel size = 1 × 0.98 × 0.98 mm.

We used voxel-based morphology (VBM) as an automated user-independent voxel-wise measurement of the associations between regional brain volumes and individual differences in reward reactivity. The VBM approach has been extensively cross-validated with manual




Fig. 1 Results of a whole-brain analysis reveal a cluster of voxels correlating with right ventral striatal activation to monetary rewards relative to losses within the midbrain which appears to encompass portions of the periaqueductal gray and ventral tegmental area (*left*). Scatterplot of extracted midbrain gray matter volume and functional win > loss reactivity from fMRI task indicates that for both the left

and right ventral striatum as midbrain volume increases, ventral striatal activation increases (*right*). These correlations remain significant when potential outliers with the highest ventral striatal BOLD reactivity are excluded (*left* $r = 0.27$, $p < 0.05$ and *right* $r = 0.58$, $p < 0.001$)

volumetric analysis (Woermann et al. 1999). The technique derives volume measurements from transformation of individual structural MR volume images into a common stereotactic space, which allows the testing of differences in sub-volumes of distinct brain regions using general linear model statistics. The VBM methodology used in this report is similar to the procedures described previously (Ashburner and Friston 2000). First, we manually adjusted volume brain images to a common orientation (origin at the anterior commissure) before all images were pre-processed using standard VBM procedures in SPM8 (<http://www.fil.ion.ucl.ac.uk/spm>). We visually examined the T_1 -weighted MPRAGE images for artifacts and structural abnormalities. We then segmented these images into gray matter, white matter and cerebrospinal fluid, after which they were again visually inspected. Gray and white matter images were normalized to standard gray matter templates in SPM8 (Ashburner and Friston 2000). Lastly, tissue probability maps were obtained by averaging across participant data, using an 8-mm FWHM Gaussian smoothing kernel. Measures of total gray matter volume were obtained from summed global signal of segmented images of gray matter.

In our earlier study, right ventral striatal activation was both the global maxima from the fMRI analysis and the best predictor of FN amplitude (Carlson et al. 2011). Therefore, participant's extracted right ventral striatal BOLD activity was used as a predictor variable with gray matter volume as the dependent variable using multiple regression within SPM8. We specifically tested whether or not increased gray matter volume would correlate with

greater BOLD activity. Age and whole-brain gray matter volume were included as covariates to control for their potentially confounding effects on regional gray matter (Ge et al. 2002). An initial whole-brain threshold was set to $P_{\text{single-tailed}} < 0.001$ and a whole-brain cluster-level family-wise error correction (FWE) was applied. Follow-up partial correlations (controlling for age and whole-brain gray matter) were performed on gray matter volume extracted from a region in the midbrain identified in the SPM8 regression.

Results

As can be seen in Fig. 1, a cluster of voxels extending bilaterally within the midbrain of the brainstem, which appears to encompass portions of the periaqueductal gray and ventral tegmental area, correlated with participant's right ventral striatal BOLD activity, $k = 165$, peak voxel at $-4, -28, -16$; $t_{38} = 5.05$, $P_{\text{FWE-corrected}} < 0.05$. No other regions correlated with right ventral striatal BOLD activity. We then performed additional partial correlations (controlling for age and whole-brain gray matter) on gray matter values extracted from the midbrain cluster identified in the SPM analyses. These follow-up analyses indicate that the association between gray matter volume and BOLD activity is present for both the left ($r = 0.32$, $P_{\text{one-tailed}} = 0.02$) and the right ($r = 0.59$, $P_{\text{one-tailed}} = 0.00003$) ventral striatum as well as the mPFC ($r = 0.38$, $P_{\text{one-tailed}} = 0.007$). In addition, midbrain gray matter volume correlated with

FN_{PCA} amplitude, $r = 0.30$, $P_{\text{one-tailed}} = 0.03$ (partial correlation controlling for age and whole-brain gray matter).

Given that FN_{PCA} amplitude has been linked to underlying VS and mPFC activation (Carlson et al. 2011), we then ran a mediation analysis to test whether the association between FN_{PCA} amplitude and midbrain gray matter was accounted for by functional activation in the right VS and mPFC. Controlling for age and whole-brain gray matter, midbrain gray matter significantly predicted functional right VS activation ($\beta = 0.73$, $P_{\text{one-tailed}} = 0.00003$). When both midbrain gray matter and functional VS activation were added as simultaneous predictors of FN_{PCA} amplitude, right VS activation continued to predict FN_{PCA} amplitude ($\beta = 0.62$, $P_{\text{one-tailed}} = 0.0003$), but midbrain gray matter did not ($\beta = -0.08$, $P_{\text{one-tailed}} = 0.34$). The Sobel test revealed that this mediation effect was statistically significant ($z = 2.91$, $P_{\text{one-tailed}} = 0.002$), indicating that the association between midbrain gray matter and FN_{PCA} amplitude was fully mediated by functional VS activation.

By contrast, mPFC activation did not mediate the relationship between midbrain gray matter and FN_{PCA} amplitude. Controlling for age and whole-brain gray matter, midbrain gray matter significantly predicted functional mPFC activation ($\beta = 0.48$, $P_{\text{one-tailed}} = 0.007$). When both midbrain gray matter and functional mPFC activation were added as simultaneous predictors of FN amplitude, however, neither association was significant (midbrain: $\beta = 0.28$, $P_{\text{one-tailed}} = 0.09$; mPFC: $\beta = 0.17$, $P_{\text{one-tailed}} = 0.16$); the Sobel test yielded no significant effect of mediation ($z = 0.95$, $P_{\text{one-tailed}} = 0.17$).

Discussion

We found that as midbrain volumes increased, fMRI reward reactivity in the VS and mPFC also increased. In addition, increases in midbrain volume were associated with an increased FN amplitude to reward. The association between midbrain volume and FN amplitude was mediated by VS reward reactivity. On the other hand, the relationship between midbrain volume and FN amplitude was not mediated by mPFC reward reactivity.

Gray matter volume is thought to represent the density of neuronal and glial cell bodies in addition to the number of dendritic branches and short range axons (Zatorre et al. 2012). Therefore, gray matter volume can be thought as a measure of local network integrity. Although the exact functional significance of this local network integrity remains unclear, presumably greater integrity in dopamine-producing neurons in the midbrain should coincide with greater dopamine production and availability in target regions such as the VS and mPFC. It has been proposed

that dysfunction specifically in the midbrain-VS circuit may be a primary etiological factor in depression (Nestler and Carlezon 2006; Russo and Nestler 2013). Indeed, recent animal data indicate that phasic dysregulation of midbrain neurons projecting to the VS—but not the mPFC—mediates depressive behavior under conditions of environmental stress (Chaudhury et al. 2013). In humans, depression is associated with lower midbrain volumes (Lee et al. 2011). Based on the present data, this structural abnormality in the midbrain may relate to functional abnormalities in depression—particularly blunted VS reactivity (Keedwell et al. 2005; Wacker et al. 2009) and FN amplitude (Bress et al. 2013a, b, 2012; Foti et al. 2011) to reward. Similarly, blunted VS activity to reward in other patient populations such as attention deficit hyperactivity disorder (Plichta and Scheres 2013) may also be linked to lower midbrain volumes. On the other end of the spectrum, greater midbrain volume may be related to over activation of the VS-mPFC reward circuit and the corresponding hypersensitivity to reward observed in pathological gamblers. Pathological gamblers have been shown to have increased midbrain activity to near miss losses in a slot machine gambling task and activation of the midbrain in these individuals correlates with activity in the VS (Habit and Dixon 2010). Based on the current results, we would expect this hypersensitivity to reward to be driven by enlarged midbrain volumes. Future research should test this possibility directly by examining changes in the structure–function relationship of reward processing within various patient populations along the reward sensitivity spectrum.

It should be noted that it is unclear from the current results as to whether lower midbrain volumes are a risk factor for blunted reward or rather a consequence of this behavior. The predominant theory for reduced gray matter volumes in depression claims that such reductions are linked to stress-mediated cell death by glucocorticoids (Woolley et al. 1990; Sheline et al. 2002). Although this mechanism has particularly been linked to reduced hippocampal and medial prefrontal cortical volumes (Bora et al. 2012; Kempton et al. 2011), it could also explain reduced midbrain gray matter in those with blunted reactivity to reward. On the other hand, decreased reactivity to reward may, over time, result in lower levels of gray matter within the dopaminergic midbrain diffuse modulatory centers. That is, it is unclear within our sample as to whether reduced gray matter within the midbrain is the cause or the effect of blunted reward reactivity. In either case, midbrain gray volumes are closely linked to current levels of reward reactivity.

Recent research has shown that learning a new skill or training results in neuroplasticity, which is observable in the MRI signal (Zatorre et al. 2012). It has recently been suggested that tracking gray matter changes related to treatments of anxiety disorders such as attention bias

modification may be useful in assessing the efficacy of such treatments (Carlson et al. 2013). That is, the relative success of treatment in reducing one's attentional bias to threat and level of anxiety should correspond to the degree of structural change within a particular brain network (i.e., the amygdala—anterior cingulate network for attention bias (Carlson et al. 2012, 2013). Similarly, given the association between functional reward reactivity and mid-brain gray matter volume observed here and the link between functional reward reactivity and symptoms of anhedonia (Keedwell et al. 2005; Wacker et al. 2009), the efficacy of treatments aimed at alleviating this symptom may be linked to structural changes within the midbrain as well as functional measures of reward reactivity. Future research should assess the extent to which successful treatment results in structural changes within the midbrain and the degree to which these potential changes are reflected in functional measures such as the FN. If structural change is predictive of treatment outcome and can be tracked through the FN, this would allow for easier implementation of tracking treatment status through physiological measures as EEG is considerably less expensive than MRI.

In sum, variability in midbrain gray matter volume accounts for variability in VS reactivity and FN amplitude to rewards. Structural abnormalities in the midbrain may underlie reward-related psychopathologies and the treatment of these disorders.

References

- Ashburner J, Friston KJ (2000) Voxel-based morphometry—the methods. *Neuroimage* 11(6):805–821. doi:10.1006/Nimg.0582
- Bora E, Fornito A, Pantelis C, Yucel M (2012) Gray matter abnormalities in major depressive disorder: a meta-analysis of voxel based morphometry studies. *J Affect Disord* 138(1–2):9–18. doi:10.1016/j.jad.2011.03.049
- Breiter HC, Gollub RL, Weisskoff RM, Kennedy DN, Makris N, Berke JD, Goodman JM, Kantor HL, Gastfriend DR, Riorden JP, Mathew RT, Rosen BR, Hyman SE (1997) Acute effects of cocaine on human brain activity and emotion. *Neuron* 19(3):591–611
- Bress JN, Smith E, Foti D, Klein DN, Hajcak G (2012) Neural response to reward and depressive symptoms in late childhood to early adolescence. *Biol Psychol* 89(1):156–162. doi:10.1016/j.biopsycho.2011.10.004
- Bress JN, Foti D, Kotov R, Klein DN, Hajcak G (2013a) Blunted neural response to rewards prospectively predicts depression in adolescent girls. *Psychophysiology* 50(1):74–81. doi:10.1111/j.1469-8986.2012.01485.x
- Bress JN, Meyer A, Hajcak G (2013b) Differentiating anxiety and depression in children and adolescents: evidence from event-related brain potentials. *J Clin Child Adolesc Psychol*. doi:10.1080/15374416.2013.814544
- Carlson JM, Beacher F, Reinke KS, Habib R, Harmon-Jones E, Mujica-Parodi LR, Hajcak G (2012) Nonconscious attention bias to threat is correlated with anterior cingulate cortex gray matter volume: a voxel-based morphometry result and replication. *Neuroimage* 59(2):1713–1718. doi:10.1016/J.Neuroimage.09.040
- Carlson JM, Foti D, Mujica-Parodi LR, Harmon-Jones E, Hajcak G (2011) Ventral striatal and medial prefrontal BOLD activation is correlated with reward-related electrocortical activity: a combined ERP and fMRI study. *Neuroimage* 57(4):1608–1616. doi:10.1016/J.Neuroimage.05.037
- Carlson JM, Cha J, Harmon-Jones E, Mujica-Parodi LR, Hajcak G (2013) Influence of the BDNF genotype on amygdalo-prefrontal white matter microstructure is linked to nonconscious attention bias to threat. *Cereb Cortex*. doi:10.1093/cercor/bht089
- Chaudhury D, Walsh JJ, Friedman AK, Juarez B, Ku SM, Koo JW, Ferguson D, Tsai HC, Pomeranz L, Christoffel DJ, Nectow AR, Ekstrand M, Domingos A, Mazei-Robison MS, Mouzon E, Lobo MK, Neve RL, Friedman JM, Russo SJ, Deisseroth K, Nestler EJ, Han MH (2013) Rapid regulation of depression-related behaviours by control of midbrain dopamine neurons. *Nature* 493(7433):532–536. doi:10.1038/nature11713
- Drevets WC, Gautier C, Price JC, Kupfer DJ, Kinahan PE, Grace AA, Price JL, Mathis CA (2001) Amphetamine-induced dopamine release in human ventral striatum correlates with euphoria. *Biol Psychiatry* 49(2):81–96
- Epstein J, Pan H, Kocsis JH, Yang Y, Butler T, Chusid J, Hochberg H, Murrough J, Strohmayr E, Stern E, Silbersweig DA (2006) Lack of ventral striatal response to positive stimuli in depressed versus normal subjects. *Am J Psychiatry* 163(10):1784–1790. doi:10.1176/appi.ajp.163.10.1784
- Fitzgerald PB, Laird AR, Maller J, Daskalakis ZJ (2008) A meta-analytic study of changes in brain activation in depression. *Hum Brain Mapp* 29(6):683–695. doi:10.1002/hbm.20426
- Foti D, Hajcak G (2009) Depression and reduced sensitivity to non-rewards versus rewards: evidence from event-related potentials. *Biol Psychol* 81(1):1–8
- Foti D, Weinberg A, Dien J, Hajcak G (2011) Event-related potential activity in the basal ganglia differentiates rewards from nonrewards: temporospatial principal components analysis and source localization of the feedback negativity. *Hum Brain Mapp* 32(12):2207–2216. doi:10.1002/hbm.21182
- Ge Y, Grossman RI, Babb JS, Rabin ML, Mannon LJ, Kolson DL (2002) Age-related total gray matter and white matter changes in normal adult brain. Part I: volumetric MR imaging analysis. *AJNR Am J Neuroradiol* 23(8):1327–1333
- Gehring WJ, Willoughby AR (2002) The medial frontal cortex and the rapid processing of monetary gains and losses. *Science* 295(5563):2279–2282. doi:10.1126/science.1066893295/5563/2279
- Habib R, Dixon MR (2010) Neurobehavioral evidence for the “Near-Miss” effect in pathological gamblers. *J Exp Anal Behav* 93(3):313–328. doi:10.1901/jeab.2010.93-313
- Hajcak G, Moser JS, Holroyd CB, Simons RF (2006) The feedback-related negativity reflects the binary evaluation of good versus bad outcomes. *Biol Psychol* 71(2):148–154. doi: 10.1016/j.biopsycho.2005.04.001
- Holroyd CB, Coles MG (2002) The neural basis of human error processing: reinforcement learning, dopamine, and the error-related negativity. *Psychol Rev* 109(4):679–709
- Holroyd CB, Larsen JT, Cohen JD (2004) Context dependence of the event-related brain potential associated with reward and punishment. *Psychophysiology* 41(2):245–253. doi:10.1111/j.1469-8986.2004.00152.x
- Holroyd CB, Hajcak G, Larsen JT (2006) The good, the bad and the neutral: electrophysiological responses to feedback stimuli. *Brain Res* 1105(1):93–101. doi:10.1016/j.brainres.2005.12.015
- Holroyd CB, Pakzad-Vaezi KL, Krigolson OE (2008) The feedback correct-related positivity: sensitivity of the event-related brain

- potential to unexpected positive feedback. *Psychophysiology* 45(5):688–697. doi:[10.1111/j.1469-8986.2008.00668.x](https://doi.org/10.1111/j.1469-8986.2008.00668.x)
- Keedwell PA, Andrew C, Williams SC, Brammer MJ, Phillips ML (2005) The neural correlates of anhedonia in major depressive disorder. *Biol Psychiatry* 58(11):843–853. doi:[10.1016/j.biopsych.2005.05.019](https://doi.org/10.1016/j.biopsych.2005.05.019)
- Kempton MJ, Salvador Z, Munafo MR, Geddes JR, Simmons A, Frangou S, Williams SC (2011) Structural neuroimaging studies in major depressive disorder. Meta-analysis and comparison with bipolar disorder. *Arch Gen Psychiatry* 68(7):675–690. doi:[10.1001/archgenpsychiatry.2011.60](https://doi.org/10.1001/archgenpsychiatry.2011.60)
- Knutson B, Bossaerts P (2007) Neural antecedents of financial decisions. *J Neurosci* 27(31):8174–8177. doi:[10.1523/JNEUROSCI.1564-07.2007](https://doi.org/10.1523/JNEUROSCI.1564-07.2007)
- Lee HY, Tae WS, Yoon HK, Lee BT, Paik JW, Son KR, Oh YW, Lee MS, Ham BJ (2011) Demonstration of decreased gray matter concentration in the midbrain encompassing the dorsal raphe nucleus and the limbic subcortical regions in major depressive disorder: an optimized voxel-based morphometry study. *J Affect Disord* 133(1–2):128–136. doi:[10.1016/j.jad.2011.04.006](https://doi.org/10.1016/j.jad.2011.04.006)
- Miltner WHR, Braun CH, Coles MGH (1997) Event-related brain potentials following incorrect feedback in a time-estimation task: evidence for a generic neural system for error detection. *J Cogn Neurosci* 9(6):788–798
- Nestler EJ, Carlezon WA Jr (2006) The mesolimbic dopamine reward circuit in depression. *Biol Psychiatry* 59(12):1151–1159. doi:[10.1016/j.biopsych.2005.09.018](https://doi.org/10.1016/j.biopsych.2005.09.018)
- O'Doherty JP, Deichmann R, Critchley HD, Dolan RJ (2002) Neural responses during anticipation of a primary taste reward. *Neuron* 33(5):815–82
- Olds J, Milner P (1954) Positive reinforcement produced by electrical stimulation of septal area and other regions of rat brain. *J Comp Physiol Psychol* 47(6):419–427
- Pizzagalli DA, Holmes AJ, Dillon DG, Goetz EL, Birk JL, Bogdan R, Dougherty DD, Iosifescu DV, Rauch SL, Fava M (2009) Reduced caudate and nucleus accumbens response to rewards in unmedicated individuals with major depressive disorder. *Am J Psychiatry* 166(6):702–710. doi:[10.1176/appi.ajp.2008.08081201](https://doi.org/10.1176/appi.ajp.2008.08081201)
- Plichta MM, Scheres A (2013) Ventral-striatal responsiveness during reward anticipation in ADHD and its relation to trait impulsivity in the healthy population: a meta-analytic review of the fMRI literature. *Neurosci Biobehav Rev*. doi:[10.1016/j.neubiorev.2013.07.012](https://doi.org/10.1016/j.neubiorev.2013.07.012)
- Plichta MM, Wolf I, Hohmann S, Baumeister S, Boecker R, Schwarz AJ, Zangl M, Mier D, Diener C, Meyer P, Holz N, Ruf M, Gerchen MF, Bernal-Casas D, Kolev V, Yordanova J, Flor H, Laucht M, Banaschewski T, Kirsch P, Meyer-Lindenberg A, Brandeis D (2013) Simultaneous EEG and fMRI reveals a causally connected subcortical-cortical network during reward anticipation. *J Neurosci* 33(36):14526–14533. doi:[10.1523/JNEUROSCI.0631-13.2013](https://doi.org/10.1523/JNEUROSCI.0631-13.2013)
- Russo SJ, Nestler EJ (2013) The brain reward circuitry in mood disorders. *Nat Rev Neurosci* 14(9):609–625. doi:[10.1038/nrn3381](https://doi.org/10.1038/nrn3381)
- Sabatinelli D, Bradley MM, Lang PJ, Costa VD, Versace F (2007) Pleasure rather than salience activates human nucleus accumbens and medial prefrontal cortex. *J Neurophysiol* 98(3):1374–1379. doi:[10.1152/jn.00230.2007](https://doi.org/10.1152/jn.00230.2007)
- Schlaepfer TE, Cohen MX, Frick C, Kosel M, Brodesser D, Axmacher N, Joe AY, Kreft M, Lenartz D, Sturm V (2008) Deep brain stimulation to reward circuitry alleviates anhedonia in refractory major depression. *Neuropsychopharmacol* 33(2):368–377. doi:[10.1038/sj.npp.1301408](https://doi.org/10.1038/sj.npp.1301408)
- Senior C (2003) Beauty in the brain of the beholder. *Neuron* 38(4):525–528
- Sheline YI, Mittler BL, Mintun MA (2002) The hippocampus and depression. *Eur Psychiatry* 17(Suppl 3):300–305
- Wacker J, Dillon DG, Pizzagalli DA (2009) The role of the nucleus accumbens and rostral anterior cingulate cortex in anhedonia: integration of resting EEG, fMRI, and volumetric techniques. *Neuroimage* 46(1):327–337. doi:[10.1016/j.neuroimage.2009.01.058](https://doi.org/10.1016/j.neuroimage.2009.01.058)
- Walter M, Bermpohl F, Mouras H, Schiltz K, Tempelmann C, Rotte M, Heinze HJ, Bogerts B, Northoff G (2008) Distinguishing specific sexual and general emotional effects in fMRI-subcortical and cortical arousal during erotic picture viewing. *Neuroimage* 40(4):1482–1494. doi:[10.1016/j.neuroimage.2008.01.040](https://doi.org/10.1016/j.neuroimage.2008.01.040)
- Wise RA (1996) Addictive drugs and brain stimulation reward. *Annu Rev Neurosci* 19:319–340. doi:[10.1146/annurev.ne.19.030196.001535](https://doi.org/10.1146/annurev.ne.19.030196.001535)
- Woermann FG, Free SL, Koepp MJ, Ashburner J, Duncan JS (1999) Voxel-by-voxel comparison of automatically segmented cerebral gray matter - A rater-independent comparison of structural MRI in patients with epilepsy. *Neuroimage* 10(4):373–384
- Woolley CS, Gould E, McEwen BS (1990) Exposure to excess glucocorticoids alters dendritic morphology of adult hippocampal pyramidal neurons. *Brain Res* 531(1–2):225–231
- Zatorre RJ, Fields RD, Johansen-Berg H (2012) Plasticity in gray and white: neuroimaging changes in brain structure during learning. *Nat Neurosci* 15(4):528–536. doi:[10.1038/nn.3045](https://doi.org/10.1038/nn.3045)
- Zink CF, Tong Y, Chen Q, Bassett DS, Stein JL, Meyer-Lindenberg A (2008) Know your place: neural processing of social hierarchy in humans. *Neuron* 58(2):273–283. doi:[10.1016/j.neuron.2008.01.025](https://doi.org/10.1016/j.neuron.2008.01.025)