

2008

***"A New Species of Freshwater Crab from Western Kenya"***

Neil Cumberlidge  
*Northern Michigan University*

Michael Dobson

Follow this and additional works at: [https://commons.nmu.edu/facwork\\_journalarticles](https://commons.nmu.edu/facwork_journalarticles)



Part of the [Biology Commons](#)

---

**Recommended Citation**

Cumberlidge, N and Dobson, M., 2008. A new species of freshwater crab (Crustacea: Decapoda: Brachyura: Potamonautidae) from western Kenya. *Proceedings of the Biological Society of Washington*, 121(4): 468–474.

This Journal Article is brought to you for free and open access by the FacWorks at NMU Commons. It has been accepted for inclusion in Journal Articles by an authorized administrator of NMU Commons. For more information, please contact [kmcdonou@nmu.edu](mailto:kmcdonou@nmu.edu), [bsarjean@nmu.edu](mailto:bsarjean@nmu.edu).

**A new species of freshwater crab of the genus *Potamonautes* MacLeay, 1838, (Brachyura: Potamoidea: Potamonautidae) from the forested highlands of western Kenya, East Africa**

Neil Cumberlidge\* and Michael Dobson

(NC) Department of Biology, Northern Michigan University, Marquette, Michigan 49855-5301, U.S.A., e-mail: ncumberl@nmu.edu

(MD) Freshwater Biological Association, The Ferry Landing, Far Sawrey, Ambleside, Cumbria, LA22 0LP, UK, e-mail: mdobson@fba.org.uk

*Abstract.*—A new species of freshwater crab, *Potamonautes subukia*, from the forested highlands of western Kenya, East Africa, is described. The small body size at maturity, smooth carapace, and slim outwardly directed first gonopod clearly distinguish this species from all other congeners.

Recent ecological studies of stream communities in the highlands of western Kenya by the second author (Dobson et al. 2007) revealed the existence of a new species of freshwater crab in the genus *Potamonautes* MacLeay, 1838, which is described herein. This genus belongs to the family Potamonautidae Bott, 1970, that includes genera from continental Africa, Madagascar, and the Seychelles (Cumberlidge et al. 2008, Yeo et al. 2008). The more than 65 species of *Potamonautes* so far described are distributed throughout sub-Saharan continental Africa, along with the Nile catchment in North Africa (Bott 1955, Stewart et al. 1995, 1997a, 1997b; Daniels et al. 1998, Stewart & Cook 1998, Cumberlidge 1999, Daniels et al. 2000, 2002; Corace et al. 2001, Gouws et al. 2001, Cumberlidge et al. 2002, Cumberlidge & Vannini 2004, Reed & Cumberlidge 2004, 2006; Cumberlidge & Tavares 2006). The new species is compared with other members of the genus and is distinguished by its small size at maturity and its possession of a unique combination of characters of the adult male (first gonopod, carapace,

thoracic sternum, and chelipeds). Specimens are deposited in the museum of the Department of Biology, Northern Michigan University, Marquette, Michigan, U.S.A. (NMU), and in The Natural History Museum, London, UK (BMNH).

*Abbreviations.*—Terminology is adapted from Cumberlidge (1999), and the classification used here follows that of Cumberlidge et al. (2008) and Ng et al. (2008). Abbreviations: cw, distance across the carapace at the widest point; cl, carapace length measured along the median line, from the frontal margin to the posterior margin; ch, carapace height (the maximum height of the cephalothorax); fw, front width measured along the frontal margin; s, thoracic sternite; s4/s5, s5/s6, s6/s7, s7/s8, sternal sulci between adjacent thoracic sternites; e, thoracic episternite; s4/e4, s5/e5, s6/e6, s7/e7, episternal sulci between adjacent thoracic sternites and episternites; a, abdominal segment; p1–p5, pereopods 1–5; asl, above sea level. All measurements are given in mm. SMNH, Swedish Museum of Natural History (Naturhistoriska riksmuseet), Stockholm, Sweden; MCZ, Museum of Comparative Zoology, Harvard University, MA, USA.

\*1 Corresponding author.




Fig. 1. *Potamonautes subukia*. Male holotype, cw 22.5 mm, from Subukia, western Kenya, NMU 18.X.2003. A, carapace, dorsal view; B, cephalothorax, frontal view; C, right cheliped, frontal view; D, left cheliped, frontal view; E, carpus and merus of right cheliped, dorsal view; F, carpus and merus of right cheliped, inferior view. Scale bar = 4.6 mm.

Superfamily Potamoidea Ortmann, 1896

Family Potamonautidae Bott, 1970

*Potamonautes subukia*, new species

Figs. 1, 2

*Material examined*.—Holotype: adult male (cw 24 mm) Kenya, Chinga River ( $0^{\circ}00'15.8''N$ ,  $36^{\circ}15'33.1''E$ ; elevation 2180 m), a small tributary of the Subukia River, Rift Valley Province, hand caught approximately 1 km southeast (upstream) of Nakuru-Nyahururu road in forest,

coll. M. Dobson, 18 Oct 2003 (NMU 18.10.2003). Paratypes: Adult male (cw 18 mm), adult female (cw 22 mm), Kakamega-Kaimosi area, western Kenya, with *P. loveni*, coll. J. McMahon, 12 Jul 1950 (BMNH 1950.12.7.21); Marun River at Kaibuibich, Sigor, Cherangani Mts. ( $1^{\circ}34'0''N$ ,  $35^{\circ}31'0''E$ ), coll. T. R. Williams, 1962 (NMU EA62.147).

*Diagnosis*.—Carapace high (ch/fw 1.3), epigastric crests distinct, postorbital crests faint, not meeting epigastric crests or

anterolateral margins; ischium of third maxilliped with either smooth or faint vertical sulcus; thoracic sternal sulcus s3/s4 reduced to two short side notches; terminal article of gonopod 1 long, angled outward at 45° angle to longitudinal axis of gonopod; broadened in mid-section (medial fold slightly higher than lateral fold); distal margin of subterminal segment indented in middle, highest on medial side lowest on lateral side; dorsal membrane very broad, widest medially, narrowest laterally; basal margins of subterminal segment of gonopod 1 with long setae.

*Description.*—Based on holotype, adult male. Carapace oval (Fig. 1A), wide (cw/fw 3.2), medium height (ch/fw 1.3); surface completely smooth semi-circular, urogastric, transverse branchial grooves faint. Front straight, relatively narrow (Fig. 1B), about one-third carapace width (fw/cw 0.32), anterior margin sharply deflexed. Postfrontal crest faint, epigastric crests distinct, postorbital crests faint, not meeting epigastric crests or anterolateral margins. Anterolateral margin between exorbital, epibranchial teeth smooth, lacking intermediate tooth; exorbital, epibranchial teeth reduced to small granules, anterolateral margin completely smooth, continuous with posterolateral margin; suborbital margin raised, completely smooth. Carapace sidewall completely smooth, divided into three parts by longitudinal (epimeral) suture (dividing suborbital, subhepatic regions from pterygostomial region), and by vertical (pleural) groove (dividing suborbital from subhepatic regions). Second sulcus s2/s3 deep, running horizontally across sternum (Fig. 2B); third sulcus s3/s4 lacking, except for two faint notches on sides; episternal sulci s4/e4, s5/e5, s6/e6, s7/e7 absent. Third maxillipeds (Fig. 2A) filling entire oral field, except for transversely oval respiratory openings at superior lateral corners; long flagellum on exopod of third maxilliped, ischium of third

maxilliped either smooth or with very faint vertical sulcus. Epistomial tooth prominent, smooth, triangular. Mandibular palp two-segmented; terminal segment single, undivided, with setae (but no hard flap) at junction between segments. Adult male abdomen slim, triangular, segments a1–a6 of male abdomen four-sided, telson (a7) a broad triangle with rounded apex; segments a5–a6 broadest. Dactylus of major cheliped broad, slightly arched, with large teeth, enclosing long thin interspace when fingers closed (Fig. 1C–F); first carpal tooth on carpus of cheliped large, pointed; second carpal tooth small, pointed, followed by several small granules; ventral margins of merus of pereopod 1 granulated; distal meral tooth distinct, pointed; superior surface of merus smooth. Pereiopods p2–p5 slender, length normal, neither elongated or shortened, p3 longest, p5 shortest, dactyli of p2–p5 tapering to point, each bearing four rows of downward-pointing short, sharp spines. Terminal article of gonopod 1 long, angled outward at 45° angle to longitudinal axis of gonopod (Fig. 2C, D); broadened in midsection (medial fold slightly higher than lateral fold); distal margin of subterminal segment indented in middle, highest on medial side lowest on lateral side; dorsal membrane very broad, widest medially, narrowest laterally; basal margins of subterminal segment of gonopod 1 with long setae.

*Size.*—A small species, the largest known specimen is the male holotype, cw 28 mm. Adults as judged by size at pubertal molt beginning around cw 16.5 mm and above (Dobson et al. 2007).

*Live coloration.*—Carapace uniformly brown, chelipeds and walking legs light brown, sternum and undersides of pereopods p1–p5 pale brown-yellow.

*Distribution.*—*Potamonautes subukia* is known from the forested highlands of western Kenya including Subukia, Kakamega, and the Cherangani Mountains.




Fig. 2. *Potamonautes subukia*. Male holotype, cw 22.5 mm, from the Subukia, western Kenya, NMU 18.X.2003. A, left third maxilliped; B, sternum; C, left first gonopod ventral view; D, left first gonopod dorsal view. Scale bar = 4.6 mm (A, B), 2.0 (C, D).

*Type locality.*—Chinga River near Subukia (0°0'15.8" N, 36°15'33.1" E), Central Province, Kenya.

*Remarks.*—*Potamonautes subukia* is closest to *P. loveni* (Colosi, 1924), a medium-sized species of freshwater crab that is also found in this part of western Kenya and eastern Uganda (Williams 1991, Cumberlidge 1997, 1998). Unfortunately, the male type of *Potamon* (*Geothelphusa*) *loveni* (SMNH 1291) from Mt Elgon, Uganda is a subadult specimen (cw 26.3 mm) whose gonopods and chelipeds had yet to attain the adult form. The carapace features of *P. subukia* were compared with those of the sub-adult type of *P. loveni*. However, in order to compare an adult male of *P. subukia* with an adult male of *P. loveni* it was necessary to examine the holotype (cw 42.6 mm) of *P. (P.) harvardi* Rathbun, 1935 (MCZ 8241), also from Mt Elgon, which is a junior synonym of *P. (G.) loveni* (Williams 1991, Cumberlidge 1997, 1998). This specimen of *P. (P.) harvardi* is an adult whose gonopods and chelipeds have reached their final form.

*Potamonautes subukia* and *P. loveni* are similar in that both species have a completely smooth carapace with a faint, almost absent postfrontal crest, both have exorbital and epibranchial teeth that are reduced to small granules, and the anterolateral margins of the carapace of both species are completely smooth, lacking teeth of any kind. However, there are a number of other characters that distinguish these two taxa. For example, the terminal article of gonopod 1 of *P. subukia* is angled outward at a 45° angle to the longitudinal axis of the gonopod, whereas the terminal article of gonopod 1 of *P. loveni* is straight and is not significantly angled outward; the first carpal tooth on the carpus of the cheliped of *P. subukia* is large and pointed, whereas this tooth is smooth and low in *P. loveni*; the distal meral tooth on the merus of the cheliped of *P. subukia* is

distinct and pointed, whereas this tooth is low and obscure in *P. loveni*; the dactylus of the cheliped of *P. subukia* has large teeth and forms a low arch that encloses a long thin interspace when the fingers are closed, whereas the dactylus of the cheliped of *P. loveni* is slim, lacks teeth, and forms a dramatic high arch that encloses a round interspace when the fingers are closed. Finally, *P. loveni* (whose adult size range is from cw 35 to 45 mm) is a much larger species than *P. subukia*: the latter species is adult at cw 16.5 mm, whereas a specimen of *P. loveni* of the same size would only be a subadult.

*Ecology.*—This species lives in streams draining the forested highlands of western Kenya and is sympatric with *P. loveni*. The life history and ecology of *P. subukia* has been studied in some detail at the type locality (Dobson et al. 2007), where it is the only crab species present. Here *P. subukia* is common with a population density that apparently exceeds 20 m<sup>-2</sup>, with year round breeding and no seasonal variation in abundance. The pubertal molt of this species was estimated based on the examination of 37 adult females that were either carrying eggs or brooding young, of which the smallest had a carapace width of 16.5 mm.

*Etymology.*—The new species has been named for the type locality of Subukia, Kenya.

#### Acknowledgments

Paul F. Clark, Miranda Lowe, and Andrew Cabrinovic of the Department of Zoology, BMNH, are thanked warmly for their hospitality during visits to the museum by the first author. The first author also acknowledges the National Science Foundation (NSF grant DES 1308 417 22) for its support. We thank John Waithaka for alerting us to the large crab population at the type locality, and Dr. Adiel Magana and Professor Jude Mathooko of the Department of Biolog-



ical Sciences, Egerton University for facilitating field collections. The type specimen was collected as part of a project funded by a grant to the second author from the Natural Environment Research Council, UK (NER/B/S/2002/00517).

### Literature Cited

- Bott, R. 1955. Die Süßwasserkrabben von Afrika (Crust., Decap.) und ihre Stammesgeschichte.—*Annales du Musée du Congo belge, (Tervuren, Belgique) C-Zoologie* 3(3)3(1):209–352.
- . 1970. Betrachtungen über die Entwicklungsgeschichte der Süßwasserkrabben nach der Sammlung des Naturhistorischen Museums in Genf/Schweiz.—*Revue Suisse Zoologie* 77(2):327–344, pls. 1, 2.
- Colosi, G. 1924. *Potamonides africains* du Museum de Stockholm.—*Arkiv für Zoologie* 16:1–24.
- Corace, R. G., N. Cumberlidge, & R. Garms. 2001. A new species of freshwater crab from Rukwazi, East Africa.—*Proceedings of the Biological Society of Washington* 114:178–187.
- Cumberlidge, N. 1997. The African and Madagascan freshwater crabs in the Museum of Natural History, Vienna (Crustacea: Decapoda: Brachyura: Potamoidea).—*Annalen des Naturhistorischen Museums in Wien* 99B:571–589.
- . 1998. The African and Madagascan freshwater crabs in the Zoologische Staatssammlung, Munich (Crustacea: Decapoda: Brachyura: Potamoidea).—*Spixiana* 21(3):193–214.
- . 1999. The freshwater crabs of West Africa. Family Potamonautidae. *Faune et Flore Tropicales* 35, Institut de recherche pour le développement IRD (ex-ORSTOM), Paris, 382 pp.
- , & M. Vannini. 2004. Ecology and taxonomy of a tree living freshwater crab (Brachyura: Potamoidea: Potamonautidae) from Kenya and Tanzania, East Africa.—*Journal of Natural History* 38:681–693.
- , & M. Tavares. 2006. Remarks on the freshwater crabs of Angola, southwestern Africa, with the description of *Potamonautes kensleyi*, new species (Brachyura: Potamoidea: Potamonautidae).—*Journal of Crustacean Biology* 26:248–257.
- , P. F. Clark, & J. Baillie. 2002. A new species of freshwater crab (Brachyura: Potamoidea: Potamonautidae) from Príncipe, Gulf of Guinea, Central Africa.—*Bulletin of the British Museum of Natural History (Zoology)*, London 68(1):13–18.
- , S. R. Daniels, & R. v. Sternberg. 2008. A revision of the higher taxonomy of the Afrotropical freshwater crabs (Decapoda: Brachyura) with a discussion of their biogeography.—*Biological Journal of the Linnean Society* 93(2):399–413.
- Daniels, S. R., B. A. Stewart, & M. J. Gibbons. 1998. *Potamonautes granularis* sp. nov. (Brachyura, Potamonautidae), a new cryptic species of river crab from the Olifants River system, South Africa.—*Crustaceana* 71:885–903.
- , & L. Burmeister. 2000. Geographic patterns of genetic and morphological divergence amongst populations of a river crab (Decapoda, Potamonautidae) with the description of a new species from mountain streams in the Western Cape, South Africa.—*Zoologica Scripta* 30:181–197.
- , G. Gouws, M. Cunningham, & C. A. Matthee. 2002. Phylogenetic relationships of the southern African freshwater crab fauna (Decapoda: Potamonautidae: *Potamonautes*) derived from multiple data sets reveal biogeographic patterning.—*Molecular Phylogenetics and Evolution* 25:511–523.
- Dobson, M., A. M. Magana, J. Lancaster, & J. M. Mathooko. 2007. Aseasonality in the abundance and life history of an ecologically dominant freshwater crab in the Rift Valley, Kenya.—*Freshwater Biology* 52:215–225.
- Gouws, G., B. A. Stewart, & P. Reavell. 2001. A new species of freshwater crab (Decapoda, Potamonautidae) from the swamp forests of KwaZulu-Natal, South Africa: biochemical and morphological evidence.—*Crustaceana* 74:137–160.
- MacLeay, W. S. 1838. Brachyurous Decapod Crustacea, Illustrations of the Zoology of South Africa 5; being a Portion of the Objects of Natural History Chiefly Collected during an Expedition into the Interior of South Africa, under the Direction of Dr. Andrew Smith, in the Years 1834, 1835, and 1836; Fitted Out by “The Cape of Good Hope Association for Exploring Central Africa.” *In*: A. Smith, ed., *Illustrations of the Zoology of South Africa; Consisting Chiefly of Figures and Descriptions of the Objects of Natural History Collected During an Expedition into the Interior of South Africa, in the Years 1834, 1835, and 1836; Fitted Out by “The Cape of Good Hope Association for Exploring Central Africa”* 5, *Invertebrata* (3):53–71.

- Ng, P. K. L., D. Guinot, & P. J. F. Davie. 2008. Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world.—The Raffles Bulletin of Zoology Supplement 17:1–286.
- Ortmann, A. E. 1896. Das system der Decapoden-Krebse.—Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie de Thiere 9:409–453.
- Rathbun, M. J. 1935. Scientific Results of an Expedition to Rain Forest Regions in Eastern Africa. 2. Crustacea.—Bulletin of the Museum of Comparative Zoology at Harvard College 79:23–28.
- Reed, S. K., & N. Cumberlidge. 2004. Notes on the taxonomy of *Potamonautes obesus* (A Milne-Edwards, 1868) and *Potamonautes calcaratus* (Gordon, 1929) (Brachyura: Potamoidea: Potamonautidae) from eastern and southern Africa.—Zootaxa 418:1–20.
- , & ———. 2006. Taxonomy and biogeography of the freshwater crabs of Tanzania, East Africa (Brachyura: Potamoidea: Potamonautidae, Platythelphusidae, Deckeniidae).—Zootaxa 1262:1–139.
- Stewart, B. A. 1997a. Morphological and genetic differentiation between populations of river crabs (Decapoda: Potamonautidae) from the Western Cape, South Africa, with a taxonomic re-examination of *Gecarcinautes brincki*.—Zoological Journal of the Linnean Society 199:1–21.
- . 1997b. Biochemical and morphological evidence for a new species of river crab *Potamonautes parvispina* sp. nov. (Brachyura, Potamonautidae).—Crustaceana 70:737–753.
- , & P. A. Cook. 1998. Identification of a new species of river crab (Decapoda: Brachyura: Potamonautidae) from South Africa using morphological and genetic data.—Journal of Crustacean Biology 18:556–571.
- , M. Coke, & P. A. Cook. 1995. *Potamonautes dentatus*, new species, a fresh-water crab (Brachyura: Potamoidea: Potamonautidae) from KwaZulu-Natal, South Africa.—Journal of Crustacean Biology 15:558–568.
- Williams, T. R. 1991. Freshwater crabs and *Simulium neavei* in East Africa. III. morphological variation in *Potamonautes loveni* (Decapoda: Potamidae).—Transactions of the Royal Society of Tropical Medicine and Hygiene 85:181–188.
- Yeo, D. C. J., P. K. L. Ng, N. Cumberlidge, C. Magalhaes, S. R. Daniels, & M. Campos. 2008. A global assessment of freshwater crab diversity (Crustacea: Decapoda: Brachyura). Pp. 275–286 in E. V. Balian, C. Lèvequè, H. Segers and M. Martens, eds., Freshwater Animal Diversity Assessment.—Hydrobiologia, vol. 595.

Associate Editor: Christopher B. Boyko.