

1-2012

Glioblastoma Derived Exosomes Induce Apoptosis in Cytotoxic T Cells Through a Fas Ligand Mediated Mechanism

Keith Z. Sabin

Richard A. Rovin
Marquette General Hospital

Johnathan Lawrence
Northern Michigan University, jolawren@nmu.edu

Robert J. Belton Jr
Northern Michigan University

Robert J. Winn
Northern Michigan University

Follow this and additional works at: http://commons.nmu.edu/facwork_journalarticles



Part of the [Biology Commons](#)

Recommended Citation

Keith Z. Sabin, Richard A. Rovin, Johnathan E. Lawrence, Robert J. Belton, and Robert J. Winn. "Glioblastoma Derived Exosomes Induce Apoptosis in Cytotoxic T Cells Through a Fas Ligand Mediated Mechanism" *Neurosurgery* 71.2 (2012): E574. Available at: http://works.bepress.com/johnathan_lawrence/4

This Journal Article is brought to you for free and open access by The Commons. It has been accepted for inclusion in Journal Articles by an authorized administrator of The Commons. For more information, please contact kclumpne@nmu.edu, kmcdonou@nmu.edu, mburgmei@nmu.edu, bsarjean@nmu.edu.

Northern Michigan University

From the SelectedWorks of Johnathan Lawrence

January 2012

Glioblastoma Derived Exosomes Induce Apoptosis in Cytotoxic T Cells Through a Fas Ligand Mediated Mechanism

Contact
Author

Start Your Own
SelectedWorks

Notify Me
of New Work



Available at: http://works.bepress.com/johnathan_lawrence/4

You could be reading the full-text of this article now if you...

Become a society member

Become a subscriber

Purchase this article

If you have access to this article through your institution,
you can view this article in

OvidSP

Neurosurgery:

August 2012

doi: 10.1227/01.neu.0000417780.38009.19

CNS ORAL PRESENTATIONS: PDF Only

190 Glioblastoma Derived Exosomes Induce Apoptosis in Cytotoxic T cells Through a Fas Ligand Mediated Mechanism.

Sabin, Keith Z. BS; Rovin, Richard A. MD; Lawrence, Johnathan E. PhD; Belton, Robert J. Jr PhD; Winn, Robert J. PhD

Abstract

INTRODUCTION: Glioblastoma multiforme deploys a number of weapons to thwart the immune system. Within the tumor microenvironment, cytotoxic T cells fall victim to Fas ligand (FasL) induced apoptosis. In prostate and colorectal cancer, exosomes can mediate this FasL induced T cell apoptosis. Exosomes are tiny, membrane bound vesicles that are released from a cell. They contain functional mRNA and protein and have cell surface molecules representative of their parent cell. It is not known if GBM derived exosomes can also mediate FasL triggered apoptosis. In this study, the role of tumor derived exosomes as the delivery vehicle for FasL is explored.

METHODS: Exosomes are isolated from the T98 cell line using differential ultracentrifugation. FasL expression in the cell line and derived exosomes is determined using reverse transcription polymerase chain reaction (RT-PCR) and Western blotting. GBM derived exosomes, recombinant FasL, and exosomes treated with an anti-FasL antibody are co-cultured with Jurkat A3 T cells. Apoptosis is measured using a caspase-8 luminescent assay.

RESULTS: FasL is expressed by the T98 cell line and is present on the surface of the cells and their exosomes (Figure 1). Caspase-8 activation is seen in T cells treated with GBM derived exosomes and recombinant FasL, but not with exosomes treated with anti-FasL antibody or exosome free supernatant (Figure 2).

CONCLUSION: GBM derived exosomes induce T cell apoptosis through a FasL mediated mechanism. This method of immune suppression has not previously been described. This research opens new avenues to antagonize GBM related immune system malfunction.

Copyright (C) by the Congress of Neurological Surgeons