

5-2009

A Left Amygdala Mediated Network for Rapid Orienting to Masked Fearful Faces

Joshua M. Carlson
joshcarl@nmu.edu

Karen S. Reinke

Reza Habib

Follow this and additional works at: http://commons.nmu.edu/facwork_journalarticles

Recommended Citation

Carlson, J. M., Reinke, K. S. & Habib, R. (2009). A left amygdala mediated network for rapid orienting to masked fearful faces. *Neuropsychologia*, 47(5), 1386-9. DOI:10.1016/j.neuropsychologia.2009.01.026

This Journal Article is brought to you for free and open access by The Commons. It has been accepted for inclusion in Journal Articles by an authorized administrator of The Commons. For more information, please contact kmcdonou@nmu.edu, bsarjean@nmu.edu.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/24414072>

A left amygdala mediated network for rapid orienting to masked fearful faces

Article in *Neuropsychologia* · May 2009

DOI: 10.1016/j.neuropsychologia.2009.01.026 · Source: PubMed

CITATIONS

65

READS

108

3 authors, including:




[Joshua M Carlson](#)

Northern Michigan University

49 PUBLICATIONS 706 CITATIONS

SEE PROFILE

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.


In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Neuropsychologia

journal homepage: www.elsevier.com/locate/neuropsychologia

Note

A left amygdala mediated network for rapid orienting to masked fearful faces

Joshua M. Carlson^{a,b,c,*}, Karen S. Reinke^{a,d}, Reza Habib^{a,b}^a Memorial Hospital of Carbondale, United States^b Department of Psychology, Southern Illinois University, United States^c Department of Biomedical Engineering, Stony Brook University School of Medicine, United States^d Department of Psychology, University of Illinois Springfield, United States

ARTICLE INFO

Article history:

Received 11 July 2008

Received in revised form 6 December 2008

Accepted 19 January 2009

Available online 30 January 2009

Keywords:

Fearful faces

Spatial attention

Amygdala

Backward masking

Emotion

ABSTRACT

A rapid response to environmental threat is highly adaptive and fearful facial expressions serve as important threat cues. The biological significance of these threat cues is demonstrated by neuroimaging findings of amygdala responses to backward masked fearful faces. Additionally, behavioral dot-probe studies reveal that backward masked fearful faces modulate spatial attention. However, little is known about the behavioral impact of the amygdala sensitivity to masked fearful faces. Using a dot-probe task with event-related functional magnetic resonance imaging (fMRI), we provide the first evidence that the amygdala is involved in orienting to backward masked fearful faces. Furthermore, this spatial attention-related amygdala response was correlated with activity in the anterior cingulate, superior temporal sulcus, and lingual gyrus.

Published by Elsevier Ltd.

Fearful faces are significant biological indicators of potential threat that are detected and processed in the emotionally responsive amygdala (Adolphs, Tranel, Damasio, & Damasio, 1994). Emotional processing theories claim the amygdala automatically mediates an orienting response to both detailed and crude threat signals (LeDoux, 1998; Ohman & Mineka, 2001). Consistent with this theoretical stance, behavioral effects in dot-probe studies indicate that unmasked (Armony & Dolan, 2002; Cooper & Langton, 2006; Pourtois, Grandjean, Sander, & Vuilleumier, 2004) and masked (Carlson & Reinke, 2008; Fox, 2002; Mogg & Bradley, 1999; Mogg & Bradley, 2002) threatening faces modulate spatial attention. Additionally, backward masking neuroimaging studies report increased amygdala activity for threat-related relative to neutral faces (Liddell et al., 2005; Morris, Ohman, & Dolan, 1998; Whalen et al., 1998). Neuroimaging dot-probe studies suggest unmasked fearful faces facilitate visual processing (Pourtois et al., 2004; Pourtois, Schwartz, Seghier, Lazeyras, & Vuilleumier, 2006) and human lesion research indicates the amygdala mediates fear-related enhancements in occipital cortex (Vuilleumier, Richardson, Armony, Driver, & Dolan, 2004).

The directing or orienting of spatial attention to crude/masked threat signals is an important aspect of the fear response; however,

the neural characteristics of this behavior have yet to be assessed. Previous fMRI fearful face dot-probe studies (e.g., Pourtois et al., 2006) have not compared directed vs. undirected spatial attention and have only assessed detailed/unmasked fearful faces. Here, we tested the hypothesis that the amygdala directs spatial attention to backward masked fearful faces through a network of brain structures that include the emotion- and attention-related anterior cingulate cortex (ACC; Armony & Dolan, 2002; Bush, Luu, & Posner, 2000).

1. Methods

Twelve (seven male and five female) right handed individuals between the ages of 18 and 35 participated in the study. Potential subjects were screened for prescription and recreational drug usage, neurological and psychological histories, and for metal that could not be removed from their body. Monetary compensation was provided to subjects for participating. Subjects gave informed consent and were treated in accordance to the guidelines of the Institutional Review Board of Southern Illinois University Carbondale.

A dot-probe task (MacLeod & Mathews, 1988) was performed while brain activity was examined with event-related fMRI. Four (two male and two female) gray scale facial identities of fearful and neutral expressions (Gur et al., 2002) were used for the initial faces. A fifth neutral female face from this database was used as the mask. Stimuli were presented using the IFIS system on an MRI-compatible LCD-screen mounted to the head coil with a field of view of 7.5°. Each trial began with a 1000 ms fixation cue. Next, two faces were simultaneously presented (33 ms) to the left and right of fixation (outer edges of the faces were separated by 15° of visual angle) and immediately masked by neutral faces (100 ms). Masks were offset by approximately 1° of visual angle on the vertical Y-axis to reduce apparent motion (Liddell et al., 2005). Horizontal shifts were not used (to reduce biasing participants' attention). Masks were followed by a left visual field (LVF) or right visual field (RVF) target dot (750 ms) and a jittered (500–2000 ms) intertrial interval (Supplementary Fig. 1).

* Corresponding author at: Department of Biomedical Engineering, Health Sciences Center 18-030, SUNY Stony Brook School of Medicine, Stony Brook, NY 11794-8181, United States. Tel.: +1 631 444 9232; fax: +1 631 444 6646.

E-mail address: carlsonjm79@gmail.com (J.M. Carlson).

With an fMRI-compatible response pad, subjects used their right index finger for LVF targets and right middle finger for RVF targets. The LCD-screen and response pad were controlled via a fiber optic cable by a control-room PC equipped with E-Prime 1.1 (Psychology Software Tools, Pittsburgh, PA).

Directed spatial attention trials consisted of one fearful and one neutral initial face with the fearful face occurring equally in either the LVF or RVF. These trials were half congruent (target dot on the same side as the fearful face) and half incongruent. Undirected attention trials consisted of either both fearful (FF) or both neutral (NN) 33 ms faces. Whereas directed attention trials are thought to contain a shift in spatial attention to the location of the fearful face, undirected trials are independent of an attentional bias to one face over the other. This difference in attentional bias enabled us to assess amygdala activity for *directed vs. undirected* spatial attention. Differential amygdala activity in the cued LVF (or RVF) vs. NN contrast should reveal activity associated both with fearful face processing and spatial attention elicited by masked fearful faces. On the other hand, differential activity in the LVF (or RVF) vs. FF+NN contrast should reveal amygdala processing strictly associated with masked fearful face-elicited spatial attention and not fearful face processing per se as the total ratio of fearful and neutral faces (1:1) is held constant in this contrast.

A 1.5T Phillips whole body scanner equipped with a head coil was used to acquire T2* weighted scans with an EPI sequence sensitive to BOLD signal using the following parameters: TR=2500 ms, TE=50 ms, flip angle=90°, matrix dimensions=64 × 64, slices=26, slice thickness=5.5 mm, gap=0. Anatomical T1-weighted structural scans were also acquired. Standard preprocessing procedures were performed in SPM5, including: image realignment corrections for head movements, slice timing corrections, normalization to standard 2 mm × 2 mm × 2 mm Montreal Neurological Institute space, and spatial smoothing with a Gaussian full-width-at-half-maximum 10 mm filter. First level single subject statistical parameter maps were created for each condition using the general linear model in SPM5. A full factorial second-level model was created with six levels (LVF: congruent and incongruent, RVF: congruent and incongruent, neutral-neutral, and fearful-fearful).

2. Results

To examine the efficacy of the masking procedure, a separate sample (see [Supplementary Methods](#)) of subjects attempted to correctly identify each trial type (LVF, RVF, FF, and NN). This control task was identical to the dot-probe task with the exception that after the initial face-mask pairing, subjects attempted to identify the trial type. Fourteen out of 15 subjects failed to identify trial types above chance ([Supplementary Table 1](#)). While the majority of control subjects failed to detect the nature of the masked faces, we do not wish to claim that facial processing was subliminal for all fMRI subjects. Nonetheless, these backward masking procedures appear to be successful at restricting the processing of the initial (masked) faces.

For reaction times (RTs), a cued visual field (left, right fearful face) by target congruency (congruent, incongruent) repeated-measures analysis of variance revealed a significant two-way interaction ($F_{1,11} = 5.68, P = 0.036$): congruent trials were faster than incongruent trials in the LVF ($P = 0.006$) but did not differ in the RVF ($P = 0.589$, [Supplementary Fig. 2](#)). This interaction is consistent with previous research ([Fox, 2002](#); [Mogg & Bradley, 1999](#); [Mogg & Bradley, 2002](#)) and suggests that in the current experiment LVF, but not RVF, fearful faces captured spatial attention.

Bilateral amygdala region of interest analyses were performed in SPM5 using Masks for Regions of Interest Analysis ([Walter et al., 2003](#)) with a search volume corrected α of 0.05. The FF vs. NN comparison resulted in increased left amygdala activity ($P = 0.005$, [Fig. 1a](#)), which confirms previous findings ([Liddell et al., 2005](#); [Morris et al., 1998](#); [Pasley, Mayes, & Schultz, 2004](#); [Whalen et al., 1998](#); [Williams et al., 2006](#); [Williams, Morris, McGlone, Abbott, & Mattingley, 2004](#)) that suggest the amygdala is sensitive to crude threatening faces. However, the laterality of this amygdala response has been inconsistent (*left*: [Pasley et al., 2004](#), *right*: [Morris et al., 1998](#), and *bilateral*: [Liddell et al., 2005](#); [Whalen et al., 1998](#); [Williams et al., 2004, 2006](#)). This lateralized amygdala activity is further explored in the discussion.

To test the hypothesis that the amygdala mediates the modulation of spatial attention to masked fearful faces we separately compared the cued LVF and cued RVF to the NN and FF+NN undirected attention conditions. As presented in [Table 1](#), the results revealed increased left amygdala activity in the LVF vs. NN ($P = 0.003$, [Fig. 1b](#)) and LVF vs. FF+NN ($P = 0.019$, [Fig. 1c](#)) attention contrasts; however masked RVF fearful faces did not enhance amygdala activity. While previous research ([Carlson & Reinke, 2008](#)) suggests that perceptual inconsistencies between the initial faces and masks do not facilitate reaction times, it is unclear how these inconsistencies might effect BOLD. However, the tight coupling between behavioral and BOLD effects (both differences in LVF, but not RVF) suggests the left amygdala is associated with enhancements in spatial attention to masked fearful faces.

Left amygdala activity (active cluster from the LVF vs. FF+NN contrast) was included as a covariate in a whole brain analysis of LVF trials to assess a potential correlated network involved in directing attention to masked fearful faces. The left amygdala significantly ($P_s < 0.05_{FWE}$) covaried with the left ACC (IACC), right lingual gyrus (rLG) and superior temporal sulcus (rSTS) in addition to other limbic and frontal areas (see [Table 1](#)). Differences in correlations were assessed by using the voxel of interest box in SPM5 to extract data from the most active voxel in the rLG, IACC, and rSTS. Follow-up Fisher's z-tests indicated that activity in the left amygdala was significantly more correlated with the IACC ($z = 2.16, P = 0.016$), rLG ($z = 2.31, P = 0.011$), and rSTS ($z = 2.18, P = 0.014$) in the LVF condition (IACC $r = 0.83$, rLG $r = 0.88$, rSTS $r = 0.85$) compared to the undirected condition (IACC $r = 0.16$, rLG $r = 0.17$, rSTS $r = .18$, [Supplementary Fig. 3](#)). A behavioral LVF "attention index" (LVF congruent–incongruent RTs) was computed for each subject and was included as an additional covariate (with left amygdala activity) in the whole brain analysis of LVF trials. For LVF trials, a whole brain conjunction analysis ([Nichols, Brett, Andersson, Wager, & Poline, 2005](#)) of correlated left amygdala activity and correlated LVF attention index activity revealed that within the previously identified attention network, that the IACC ($P < 0.001_{uncorrected}$) and rLG ($P = 0.006_{uncorrected}$) covaried with both measures (see [Table 1](#) for additional areas of covariation).




Fig. 1. Left amygdala ($y = 2$) responsivity to (a) masked fearful faces, (b) LVF fearful face and attention-related processing, and (c) LVF fearful face-elicited spatial attention. Activation displayed at $P = 0.05$, adjusted for search volume.

Table 1
Masked fearful face-related spatial attention activity.

Analysis and region	Hemisphere	MNI coordinates			Voxels	Maximally activated voxel	
		x	y	z		t-value	p-value
Amygdala ROI analyses							
FF vs. NN	L	−26	0	−24	71*	2.64	0.005
LVF vs. NN	L	−30	0	−22	57*	2.87	0.003
LVF vs. NN + FF	L	−30	0	−22	17*	2.11	0.019
Whole brain left amygdala covariation (LVF trials)							
Anterior cingulate	L	−4	46	−4	329***	6.85	<0.05 _{FWE}
Lingual gyrus	R	10	−52	4	142***	8.80	<0.05 _{FWE}
Superior temporal sulcus	R	50	−32	−4	82***	7.72	<0.05 _{FWE}
Parahippocampal gyrus/hippocampus	L	−22	−18	−16	112***	7.74	<0.05 _{FWE}
Cingulate	L	−4	24	30	327***	6.78	<0.05 _{FWE}
Inferior frontal gyrus	L	−38	28	6	29***	6.76	<0.05 _{FWE}
Whole brain conjunction of left amygdala and attention index covariation (LVF trials)							
Anterior cingulate	L	−8	48	−8	52**	5.97	<0.001
Lingual gyrus	R	10	−54	2	4**	3.18	<0.006
Cingulate	L	−6	26	32	43**	8.05	<0.001
Anterior temporal pole	L	−34	8	−18	123**	5.73	<0.001
Parahippocampal gyrus/hippocampus	L	−22	−30	−14	26**	5.23	<0.001
Parahippocampal gyrus	R	26	−48	8	33**	4.97	<0.001
Brain stem	R	4	−20	−14	6**	4.40	0.001

* $p < 0.05_{\text{svc}}$.** $p < 0.01_{\text{uncorrected}}$ (conjunction).*** $p < 0.005_{\text{FDR}}$.

3. Discussion

Consistent with previous research (Carlson & Reinke, 2008; Fox, 2002; Mogg & Bradley, 1999; Mogg & Bradley, 2002), our behavioral results indicate that backward masked LVF fearful faces enhanced spatial attention. In line with these behavioral results, we found left amygdala activity for LVF fearful faces. On the other hand, RVF fearful faces did not enhance RTs nor elicit amygdala activity. Collectively, these results suggest the left amygdala mediates the facilitation of spatial attention to masked fearful faces. Additionally, left amygdala activity was found to be positively correlated with IACC, rSTS, and rLG activity when visuospatial attention was directed to backward masked LVF fearful faces, but not when attention was undirected. Finally, the results revealed that the IACC and rLG were associated both with subjects' attention-related left amygdala activity and behavioral attention index.

The observed ipsilateral amygdala processing of masked fearful faces contradicts the contralateral processing typically observed in the visual system. However, a review of the neuroimaging literature (Baas, Aleman, & Kahn, 2004) indicates a greater reported incidence of left, compared to right, amygdala activity. Indeed, we found left, but not right, amygdala activity in the FF vs. NN and LVF attention contrasts. Evidence from retrograde tracing studies in rats indicates that the amygdala receives bilateral input from visual nuclei in the brainstem, which may provide a neuroanatomical subcortical substrate for ipsilateral visual processing (Usunoff, Itzev, Rolfs, Schmitt, & Wree, 2006). The aforementioned amygdala characteristics may explain our left lateralized effects.

We provide the first evidence that the left amygdala mediates enhancements of spatial attention to masked fearful faces through a network including the IACC, rSTS, and rLG. Based on previous findings, we speculate that the amygdala detects fearful (or emotional) stimuli relayed from subcortical visual nuclei (Liddell et al., 2005; Morris, DeGelder, Weiskrantz, & Dolan, 2001; Pasley et al., 2004; Usunoff et al., 2006) and interacts with the emotion- and attention-related ACC (Armony & Dolan, 2002; Bush et al., 2000) to modulate contralateral visual processing (rLG and rSTS) through amygdala dependent feedback (Vuilleumier et al., 2004). The STS processes changeable aspects of face perception, such as eye gaze direction and emotional expression (Haxby, Hoffman, & Gobbini,

2000) and has been found to be correlated with the amygdala during crude fearful face processing (Jiang & He, 2006). Our results add to findings of unmasked fearful face-elicited attentional enhancements in visual cortex (Pourtois et al., 2004, 2006) and suggest this enhancement may be mediated by the amygdala. The amygdala and ACC may be involved in orienting spatial attention to crude threat signals, whereas enhancements in contralateral visual processing may reflect the site of attentional capture. While we did not assess the neural structures involved in the disengagement of spatial attention, previous research (with unmasked faces) points to intraparietal and orbitofrontal areas (Armony & Dolan, 2002; Pourtois et al., 2006). In sum, the threat- or emotion-initiated left amygdala, IACC and rLG network may reflect a rapid and automatic modulation of the "spotlight of attention" (Posner, 1980), which produces an adaptive facilitation in behavioral responses to stimuli within this retinotopically distinct visual location.

Acknowledgements

We would like to thank Amy Querton at the University of Illinois Springfield for assistance in data collection for the control experiment. We would also like to thank Julie Alstat and Gary Etherton from Carbondale Memorial Hospital for assisting in MRI data acquisition.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.neuropsychologia.2009.01.026.

References

- Adolphs, R., Tranel, D., Damasio, H., & Damasio, A. (1994). Impaired recognition of emotion in facial expressions following bilateral damage to the human amygdala. *Nature*, 372(6507), 669–672.
- Armony, J. L., & Dolan, R. J. (2002). Modulation of spatial attention by fear-conditioned stimuli: An event-related fMRI study. *Neuropsychologia*, 40(7), 817–826.
- Baas, D., Aleman, A., & Kahn, R. S. (2004). Lateralization of amygdala activation: a systematic review of functional neuroimaging studies. *Brain Research. Brain Research Reviews*, 45(2), 96–103.
- Bush, G., Luu, P., & Posner, M. I. (2000). Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Science*, 4(6), 215–222.

- Carlson, J. M., & Reinke, K. S. (2008). Masked fearful faces modulate the orienting of covert spatial attention. *Emotion*, 8(4), 522–529.
- Cooper, R. M., & Langton, S. R. (2006). Attentional bias to angry faces using the dot-probe task? It depends when you look for it. *Behaviour Research and Therapy*, 44(9), 1321–1329.
- Fox, E. (2002). Processing emotional facial expressions: the role of anxiety and awareness. *Cognitive, Affective & Behavioral Neuroscience*, 2(1), 52–63.
- Gur, R. C., Sara, R., Hagendoorn, M., Marom, O., Hughett, P., Macy, L., et al. (2002). A method for obtaining 3-dimensional facial expressions and its standardization for use in neurocognitive studies. *Journal of Neuroscience Methods*, 115(2), 137–143.
- Haxby, J. V., Hoffman, E. A., & Gobbini, M. I. (2000). The distributed human neural system for face perception. *Trends in Cognitive Science*, 4(6), 223–233.
- Jiang, Y., & He, S. (2006). Cortical responses to invisible faces: dissociating subsystems for facial-information processing. *Current Biology*, 16(20), 2023–2029.
- LeDoux, J. E. (1998). *The emotional brain the mysterious underpinnings of emotional life*. London: Weidenfeld and Nicholson.
- Liddell, B. J., Brown, K. J., Kemp, A. H., Barton, M. J., Das, P., Peduto, A., et al. (2005). A direct brainstem-amygdala-cortical 'alarm' system for subliminal signals of fear. *Neuroimage*, 24(1), 235–243.
- MacLeod, C., & Mathews, A. (1988). Anxiety and the allocation of attention to threat. *The Quarterly Journal of Experimental Psychology A*, 40(4), 653–670.
- Mogg, K., & Bradley, B. P. (1999). Orienting of attention to threatening facial expressions presented under conditions of restricted awareness. *Cognition and Emotion*, 13(6), 713–740.
- Mogg, K., & Bradley, B. P. (2002). Selective orienting of attention to masked threat faces in social anxiety. *Behaviour Research and Therapy*, 40(12), 1403–1414.
- Morris, J. S., DeGelder, B., Weiskrantz, L., & Dolan, R. J. (2001). Differential extrageniculostriate and amygdala responses to presentation of emotional faces in a cortically blind field. *Brain*, 124(Pt 6), 1241–1252.
- Morris, J. S., Ohman, A., & Dolan, R. J. (1998). Conscious and unconscious emotional learning in the human amygdala. *Nature*, 393(4), 467–470.
- Nichols, T., Brett, M., Andersson, J., Wager, T., & Poline, J. B. (2005). Valid conjunction inference with the minimum statistic. *Neuroimage*, 25, 653–660.
- Ohman, A., & Mineka, S. (2001). Fears, phobias, and preparedness: toward an evolved module of fear and fear learning. *Psychology Review*, 108(3), 483–522.
- Pasley, B. N., Mayes, L. C., & Schultz, R. T. (2004). Subcortical discrimination of unperceived objects during binocular rivalry. *Neuron*, 42(1), 163–172.
- Posner, M. I. (1980). Orienting of attention. *The Quarterly Journal of Experimental Psychology*, 32(1), 3–25.
- Pourtois, G., Grandjean, D., Sander, D., & Vuilleumier, P. (2004). Electrophysiological correlates of rapid spatial orienting towards fearful faces. *Cerebral Cortex*, 14(6), 619–633.
- Pourtois, G., Schwartz, S., Seghier, M. L., Lazeyras, F., & Vuilleumier, P. (2006). Neural systems for orienting attention to the location of threat signals: an event-related fMRI study. *Neuroimage*, 31(2), 920–933.
- Usunoff, K. G., Itzev, D. E., Rolfs, A., Schmitt, O., & Wree, A. (2006). Brain stem afferent connections of the amygdala in the rat with special references to a projection from the parabrachial nucleus: a fluorescent retrograde tracing study. *Anatomy and Embryology (Berlin)*, 211(5), 475–496.
- Vuilleumier, P., Richardson, M. P., Armony, J. L., Driver, J., & Dolan, R. J. (2004). Distant influences of amygdala lesion on visual cortical activation during emotional face processing. *Nature Neuroscience*, 7(11), 1271–1278.
- Walter, B., Blecker, C., Kirsch, P., Sammer, G., Schienle, A., Stark, R., et al. (2003). *MARINA: An easy to use tool for the creation of Masks for Region of Interest Analyses*. Paper presented at the 9th International Conference on Functional Mapping of the Human Brain, June 19–22, New York, NY.
- Whalen, P. J., Rauch, S. L., Etcoff, N. L., McInerney, S. C., Lee, M. B., & Jenike, M. A. (1998). Masked presentations of emotional facial expressions modulate amygdala activity without explicit knowledge. *Journal of Neuroscience*, 18(1), 411–418.
- Williams, L. M., Liddell, B. J., Kemp, A. H., Bryant, R. A., Meares, R. A., Peduto, A. S., et al. (2006). Amygdala-prefrontal dissociation of subliminal and supraliminal fear. *Human Brain Mapping*.
- Williams, M. A., Morris, A. P., McGlone, F., Abbott, D. F., & Mattingley, J. B. (2004). Amygdala responses to fearful and happy facial expressions under conditions of binocular suppression. *Journal of Neuroscience*, 24(12), 2898–2904.