

EMG INTER-LIMB ASYMMETRY IN BENCH PRESS EXERCISE IN ELITE PARALYMPICS WEIGHTLIFTERS

Esteban Aedo-Muñoz^{1,2,3,6}, Celso Sánchez-Ramírez², Christopher Moya-Jofre¹,
Alejandro Bustamante-Garrido^{1,4}, Mauricio Araya-Ibacache¹, Fabio Dal Bello³,
Ciro José Brito⁶ & Bianca Miarka^{5,6}

Biomechanics Laboratory, Chilean High Performance Center, Santiago, Chile¹
Physical Activity, Sport and Health Sciences Laboratory, Universidad de Santiago, Chile²
Physical Activity and Sports Science Master Program, Universidad Santo Tomás,
Santiago, Chile³
Kinesiology Department, Faculty of Medicine, Universidad de Chile⁴
Postgraduate Program in Physical Education, School of Physical Education and Sports,
Department of Fights. Federal University of Rio de Janeiro⁵
Postgraduate Program in Physical Education. Physical Education Department. Federal
University of Juiz de Fora, Brazil⁶

The purpose of this study was to describe EMG inter-limb asymmetry in three muscle groups in a sample of Paralympic weightlifters during an 80% RM bench press execution. The sample was composed of 7 subjects belonging to the Chilean elite powerlifting. It assesses surface electromyography activity in major pectoral, deltoideus anterior and triceps brachii. The magnitude of the response was calculated through root mean square (RMS). It was calculated Symmetry Index (SI) like interlimb differences mensure. Only the pectoralis major muscle showed significative differences between limbs (right 84.7 ± 41.3 ; left 66.1 ± 19.3 RMS) ($p=0.05$) and the SI median greatest value ($19.74 \pm 24.59\%$). Anterior deltoideus showed high individual differences in two athletes with upper 80% SI values. It is established the need to obtain more studies that assess asymmetry with the objective to decrease this injuries risk factor.

KEY WORDS: Inter-limb Asymmetry, Paralympic Biomechanics, EMG, Muscle activity.

INTRODUCTION: The selection of strength exercises represents important elements in sports training and performance. One of the most frequent and recommended exercises is the bench press exercise. This exercise is widely used by professionals of physical activity and sports to achieve strength increases in the upper limbs, both in conventional and paralympic sports (Gomo & Van Den Tillaar, 2016) (Calatayud et al., 2015) (Lehman, 2005). In the case of competitors in Paralympic weightlifting, they must first meet the minimum disability criteria in order to participate (Committee, 2015). Frequently, these athletes make changes in the bench press in terms of grip width, in addition to the levels of body inclination, which has shown certain variations in the recruitment of motor units (Hernández-Rodríguez et al., 2001) (Lehman, 2005) (Aedo-Muñoz, Herrera-Valenzuela, Bustamante-Garrido, & Letelier-Castro, 2014). Surface electromyography evaluations determine the electrical activity used by the superficial muscles, estimating the neuromuscular activity generated, in terms of the frequency and amplitude of these signals (Cifrek, Medved, Tonkovic, & Ostojic, 2009) (Basmajian & De Luca, 1985) (Konrad, 2005). Currently, several studies have characterized the electromyographic activity in the bench press exercise for different inclinations (Hernández-Rodríguez et al., 2001) (Aedo-Muñoz et al., 2014) (Padulo, Laffaye, Chaouachi, & Chamari, 2015), whereas on the other hand other authors realize it in relation to the width of the one taken (Barnett, Kippers, & Turner, 1995) (Lehman, 2005). However, the differences in neuromuscular control, measured with EMG, show a high relationship in relation to risk of injury that athletes could present during their physical preparation (Zebis et al., 2011). These asymmetries between extremities are studied even through kinetic instruments in different sports (Schiltz et al., 2009) (Andrade et al., 2013), being the electromyographic activity the measure that delivers higher level of asymmetry, even greater than the kinetic and kinematic (McAllister & Costigan, 2019). However, the evaluation of Paralympic athletes is scarce.

The purpose of this study is was to describe EMG inter-limb asymmetry in three muscle groups in a sample of Paralympic weightlifters during an 80% RM bench press execution.

METHODS:

The sample was composed of 7 elite para-powerlifting who have been members of Chilean teams for over 5 years. Throughout the study, they subjects continued with their normal daily trainings, following the regime determined by their trainer. Prior to testing, a short explanation was provided of the study, aims, procedures, and possibility of risk. This was accompanied immediately by a standardized protocol of warming up, which consisted of 10 series of bench press of 40% RM, followed by 10 minutes of stretching exercises. After the warm up, the subjects were introduced to the protocol, with an incremental intensity of; 50%-60%-70%-75% RM every five-minute without fatigue. The evaluation procedure consisted of every subject performing a sub-maximal test of 80% RM. This was considered to be the visual execution analysis and the EMG parameters. The gathered information was: surface electromyography activity in pectoralis major, anterior deltoideus and triceps brachii. The position of the electrodes was marked on the skin according to the SENIAM® (Surface Electromyography for the Non-invasive Assessment of Muscle)(Stegeman & Hermens, 2007). The electromyography signal was recorded with Delsys® Trigno, using silver surface bipolar electrodes (99%, 1 mm width and 10 mm length), with an inter-electrode distance of 10 mm. (Delsys Model Inc. Boston. M. USA). The signals registered were pre-amplific, with a common mode rejection ratio of 92dB and a gain of 1kHz (Delsys Inc. Boston. USA). The electromyography signs were processed in a macrocomputer, with IGOR PRO Wavemetrics 5.01, and were rectified completely and passed through a 6Hz low-pass digital filter. The onset muscle activation was defined as basal level, which corresponded to the average amplitude recorded in a window before activation, with a threshold corresponding to the baseline value plus 10 standard deviations from that window. When the onset of the muscular electrical activity was determined, the magnitude of the response was calculated through RMS (root mean square). To establish the EMG variation, it was not possible to perform a Maximum Voluntary Contraction (MVC) test because the sample cannot perform MVC in its sport planning model at this stage.

The descriptive statistics used were the median, major and minor values, and percentiles 25 and 75, as a measure of dispersion. The comparison between the right side and the left side of each subject was carried out using the Wilcoxon test. The "symmetry index" (SI) was calculated through the following equation:

$$SI = \frac{X_R - X_L}{\frac{1}{2} (X_R + X_L)} \times 100\%$$

Where SI = Symmetry Index, X_R = Value of the right side, X_L = V Value of the left side

SI corresponds to the percentage of asymmetry of one of the sides in relation to the other (VanZant, McPoil, & Cornwall, 2014). When SI = 0, this indicates the existence of perfect symmetry. If the value is negative, then it is assumed that the asymmetry goes to the left side, and, if it is positive, to the right side.

The respective "SI" indices were compared in the three muscle groups evaluated through ANOVA-style analysis of variance. The Bonferroni method was used as post hoc. A " $p \leq 0.05$ " was considered as a value of statistical significance.

RESULTS: Table 1 shows the descriptive data of each one of the evaluations made by the 7 athletes involved in the study. It is then possible to see that the group is heterogeneous regarding age and personal best brand. From the observation of the EMG values, it can be seen that there is an inequality between the RMS values, of the right and left sides, for the

three muscular portions evaluated in each one of the subjects. However, based on the Wilcoxon test, it was established that there were no significant differences between the right and left sides in triceps brachii ($p = 0.16$), nor in anterior deltoideus ($p = 0.94$), but they were observed in the muscle group "pectoralis major" ($p = 0.05$). While the effect sizes were performed through Cohen's d test, establishing mean values for triceps brachii ($d = 0.41$) and pectoralis major ($d = 0.58$), however in the anterior deltoid muscle the effect size is low ($d = 0.11$).

Table 1: Descriptive values of RMS in each of the muscle groups studied, for each of the subjects within the sample.

Athlete	Age (years)	PB (kg)	80% PB (kg)	Triceps Brachii (rms)		Anterior Deltoideus (rms)		Pectoralis Major (rms)	
				Right	Left	Right	Left	Right	Left
1	23	106	84.8	111.8	244.6	148.4	184.0	43.5	53.9
2	38	180	144.0	311.2	361.4	131.7	426.1	103.6	94.5
3	29	65	52.0	129.3	131.3	184.5	209.8	69.5	56.3
4	19	106	84.8	95.2	83.5	103.8	77.9	58.9	46.4
5	22	115	92.0	137.3	130.9	169.8	162.4	60.1	49.4
6	37	182	145.6	63.1	122.7	283.1	120.5	166.1	87.2
7	45	175	140.0	156.8	182.4	191.8	100.1	91.2	74.8
Minimum	19.0	65.0	52.0	63.1	83.5	103.8	77.9	43.5	46.4
P25%	22.0	106.0	84.8	95.2	122.7	131.7	100.1	58.9	49.4
Median	29.0	115.0	92.0	129.3	131.3	169.8	162.4	69.5	56.3
P75%	38.0	180.0	144.0	156.8	244.6	191.8	209.8	103.6	87.2
Maximum	45.0	182.0	145.6	311.2	361.4	283.1	426.1	166.1	94.5
				Wilcoxon	0.16	0.94	0.05		
				Cohen's d	0.41	0.11	0.58		

PB = Personal Best; RMS = root mean square; P25% = 25% percentile; P75% = 75% percentile

The median of the "SI" for each muscle group was -14.92%, 4.44% and 19.74%, for triceps brachii, anterior deltoideus and pectoralis major respectively.

Figure 1 shows the SI values obtained for each of the subjects. There it is observed that, for the triceps braquii muscle group, subjects 1 and 6 show the greatest differences (SI = -74.5% and -64.2% respectively), asymmetries oriented towards a predominance by the left side. In previous deltoideus, the greatest asymmetries were observed in subjects 2, 6 and 7, with SI values of -105.6%, 80.6% and 62.8% respectively. In the pectoralis major group, subject number 6 showed the greatest asymmetry, with SI = 62.2%, followed by subject 1 with SI = -21.3%.

From the individual analysis, it is possible to indicate that the subject No. 6 is the one who has the highest SI values and who holds the highest personal record. This tendency is repeated in subject 2, who follows him in his personal mark and who has the greatest asymmetry in the previous group deltoideus. Subjects 3, 4 and 5 presented the lowest SI values, not exceeding the 24% asymmetry value.

The analysis of the variance did not show significant differences between different SI values within the three muscle groups ($F = 1.87$, $p = 0.21$).




Figure 1: Descriptive graph of the SI values obtained for each of the subjects in the sample, in each of the muscle groups studied.

DISCUSSION: The aim of this study is to describe the inter-limbs asymmetries that the EMG signal presents in three distinctive muscle groups, in a sample of paralympics weightlifters. The sample used in this study is small to generalize its behavior, determining a starting point for future research. In this regard, it is necessary to comment that there is little officially published literature that has based its experiences on this type of sample. Additionally, to date there are no studies that have focused on the variable of asymmetry of muscle activation in the bench press exercise.

What is most relevant of the results found is the fact that the only muscle group that presents significant inter-limbs differences is the pectoralis major, which is the most requested during this exercise (Schoenfeld, Contreras, Vigotsky, & Ogborn, 2016), a result that agrees with the fact that this muscle group has the highest median SI values, close to 20% asymmetry and with predominance towards the right side. Due to the importance that this muscle group has for the execution of the aforementioned exercise, it is necessary to pay attention to the presence of these asymmetries, as it is known that this is a predisposing factor for injuries in other parts of the body (Ning, Haddad, Jin, & Mirka, 2011).

Another interesting result is that the highest absolute values for SI were recorded in the "anterior deltoideus" muscle group, with a value that exceeded 100% asymmetry towards the left side, which indicates that the left side exerts a similar EMG activity to the double of the opposite side, and in another individual with a value of 80.6% towards the right side. This result indicates a need to begin to monitor throughout the training process the different adaptations experienced by both sides of the body, in order to safeguard a correct and efficient execution of the technique (Rahnama, Reilly, Lees, & Graham-Smith, 2003). It is possible that a larger sample could register differences in those muscles that this study did not observe.

CONCLUSION: Within a sample of Paralympic athletes it was possible to establish the presence of high levels of asymmetries in the EMG activation of the muscle groups "pectoralis major" and "anterior deltoideus". The need to have more studies that tend to the evaluation of asymmetries is established, with the objective of reducing this risk factor and avoiding the appearance of injuries.

REFERENCES

- Aedo-Muñoz, E., Herrera-Valenzuela, T., Bustamante-Garrido, A., & Letelier-Castro, F. (2014). Actividad EMG del músculo pectoral mayor en los ejercicios de press banco. *Revista Akademeia*, 4.
- Andrade, M. S., Vancini, R. L., de Lira, C. A. B., Mascarin, N. C., Fachina, R. J. F. G., & da Silva, A. C. (2013). Shoulder isokinetic profile of male handball players of the Brazilian National Team. *Brazilian Journal of Physical Therapy*, 17(6), 572–578. <https://doi.org/10.1590/S1413-35552012005000125>
- Barnett, C., Kippers, V., & Turner, P. (1995). Effects of Variations of the Bench Press Exercise on the EMG Activity of Five Shoulder Muscles. *The Journal of Strength & Conditioning Research*, 9(4). Retrieved from https://journals.lww.com/nsca-jscr/Fulltext/1995/11000/Effects_of_Variations_of_the_Bench_Press_Exercise.3.aspx
- Basmajian, J., & De Luca, C. (1985). *Muscles Alive: Their Functions Revealed by Electromyography*. Williams & Wilkins. Retrieved from <https://books.google.cl/books?id=H9pqAAAAMAAJ>
- Calatayud, J., Borreani, S., Colado, J. C., Martin, F., Tella, V., & Andersen, L. L. (2015). Bench press and push-up at comparable levels of muscle activity results in similar strength gains. *Journal of Strength and Conditioning Research*, 29(1), 246–253. <https://doi.org/10.1519/JSC.0000000000000589>
- Cifrek, M., Medved, V., Tonkovic, S., & Ostojic, S. (2009). Surface EMG based muscle fatigue evaluation in biomechanics. *Clinical Biomechanics (Bristol, Avon)*, 24(4), 327–340. <https://doi.org/10.1016/j.clinbiomech.2009.01.010>
- Committee, P. (2015). *Powerlifting Classification Rules and Regulations*. Retrieved from https://www.paralympic.org/sites/default/files/document/150616104902276_IPC%2BPowerlifting%2BClassification%2BRules%2Band%2BRegulations%2Bversion%2B2015_0.pdf
- Gomo, O., & Van Den Tillaar, R. (2016). The effects of grip width on sticking region in bench press. *Journal of Sports Sciences*, 34(3), 232–238. <https://doi.org/10.1080/02640414.2015.1046395>
- Hernández-Rodríguez, R., García-Manso, M. J., Tous Fajardo, J., Ortega Santana, F., Vega Melián, F., & Gallud Marrero, I. (2001). Actividad electromiográfica del músculo pectoral mayor en los movimientos de press de banca inclinado y declinado respecto al press de banca horizontal. *Apunts. Medicina de l'Esport*, 36(136), 15–22. [https://doi.org/https://doi.org/10.1016/S1886-6581\(01\)75989-0](https://doi.org/https://doi.org/10.1016/S1886-6581(01)75989-0)
- Konrad, P. (2005). *The ABC of emg. A practical introduction to kinesiological electromyography*. Boston. <https://doi.org/10.1016/j.jacc.2008.05.066>
- Lehman, G. J. (2005). The influence of grip width and forearm pronation/supination on upper-body myoelectric activity during the flat bench press. *Journal of Strength and Conditioning Research*, 19(3), 587–591. <https://doi.org/10.1519/R-15024.1>
- McAllister, M., & Costigan, P. (2019). Evaluating movement performance: What you see isn't necessarily what you get. *Human Movement Science*, 64, 67–74. <https://doi.org/10.1016/j.humov.2019.01.003>
- Ning, X., Haddad, O., Jin, S., & Mirka, G. A. (2011). Influence of asymmetry on the flexion relaxation response of the low back musculature. *Clinical Biomechanics (Bristol, Avon)*, 26(1), 35–39. <https://doi.org/10.1016/j.clinbiomech.2010.08.012>
- Padulo, J., Laffaye, G., Chaouachi, A., & Chamari, K. (2015). Bench press exercise: the key points. *The Journal of Sports Medicine and Physical Fitness*, 55(6), 604–608.
- Rahnama, N., Reilly, T., Lees, A., & Graham-Smith, P. (2003). Muscle fatigue induced by

exercise simulating the work rate of competitive soccer. *Journal of Sports Sciences*, 21(11), 933–942. <https://doi.org/10.1080/0264041031000140428>

Schiltz, M., Lehance, C., Maquet, D., Bury, T., Crielaard, J.-M., & Croisier, J.-L. (2009). Explosive strength imbalances in professional basketball players. *Journal of Athletic Training*, 44(1), 39–47. <https://doi.org/10.4085/1062-6050-44.1.39>

Schoenfeld, B. J., Contreras, B., Vigotsky, A., & Ogborn, D. I. (2016). Upper body muscle activation during low-versus high-load resistance exercise in the bench press. *Isokinetics and Exercise Science*, 24(October 2017), 217–224. <https://doi.org/10.3233/IES-160620>

Stegeman, D., & Hermens, H. (2007). *Standards for surface electromyography: The European project Surface EMG for non-invasive assessment of muscles (SENIAM)* (Vol. 1). Retrieved from <http://www.med.uni-jena.de/motorik/pdf/stegeman.pdf>,

VanZant, R. S., McPoil, T. G., & Cornwall, M. W. (2014). Symmetry of Plantar Pressures and Vertical Forces in Healthy Subjects During Walking. *Journal of the American Podiatric Medical Association*, 91(7), 337–342. <https://doi.org/10.7547/87507315-91-7-337>

Zebis, M. K., Andersen, L. L., Pedersen, M. T., Mortensen, P., Andersen, C. H., Pedersen, M. M., ... Sjogaard, G. (2011). Implementation of neck/shoulder exercises for pain relief among industrial workers: a randomized controlled trial. *BMC Musculoskeletal Disorders*, 12, 205. <https://doi.org/10.1186/1471-2474-12-205>